

March 2026, Volume 2, Issue I.

doi: 10.65287/josta.202602.7c35

## Adoption Behaviour of Indigenous Agricultural and Ethnoveterinary Practices Among Tribal Farmers of Kalrayan Hills, Tamil Nadu

Kaviya P\*   
Annamalai University

Natarajan M   
Annamalai University

### Abstract

Indigenous agricultural and ethnoveterinary practices are integral components of the traditional knowledge systems of tribal farming communities and play a vital role in promoting sustainable agriculture and livestock health. The present study assessed the adoption behaviour of tribal farmers toward indigenous agricultural and ethnoveterinary practices in the Kalrayan Hills of Tamil Nadu, a region rich in tribal culture and traditional knowledge. An ex-post facto research design was employed, and data were collected from tribal farmers using a well-structured and pre-tested interview schedule. Adoption behaviour was analyzed across various indigenous practices and respondents were categorized based on their adoption levels. The results indicated that a majority of farmers exhibited medium to high adoption, particularly in crop protection, soil fertility management, animal healthcare and disease prevention. Factors such as age, farming experience, inheritance of traditional knowledge and access to local resources influenced adoption. Despite increased exposure to modern technologies, indigenous practices remain preferred due to their cost-effectiveness, eco-friendliness and cultural relevance. The study emphasizes the need for documentation, validation and integration of indigenous practices into formal extension programmes to strengthen sustainable agriculture and livelihood security among tribal farmers.

**Keywords:** Traditional farming systems, Indigenous technical knowledge, Animal health management, Sustainable agriculture, Tribal livelihood systems.

Received: 2026-02-05 | Accepted: 2026-03-02 | Published: 2026-03-03

© 2026 The Author(s). Licensee: Journal of Sustainable Technology in Agriculture. This is an Open Access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

## 1. Introduction

Ethno-agriculture and ethnoveterinary science encompass the study of traditional, region-specific knowledge related to the use of plants and animals by indigenous and tribal communities for agricultural and livestock management. The term “ethno” was first introduced by J. W. Harshberger in 1895 to describe the systematic study of plants and domesticated animals used by primitive and aboriginal societies (Vivekanandan 1994; Kumari *et al.* 2018). Although such knowledge systems have existed since the early stages of human civilization, ethno-agriculture and ethnoveterinary science emerged as recognized academic disciplines within environmental and agricultural sciences during the twentieth century.

Ethno-agricultural and ethnoveterinary practices are intrinsically linked to food and nutritional security, healthcare, livelihood sustenance, cultural beliefs, cottage industries, economic upliftment, biodiversity conservation and the sustainable utilization of natural resources. These practices, commonly referred to as Indigenous Technical Knowledge (ITK), reflect the deep-rooted cultural, spiritual and ecological relationships between tribal communities and their surrounding environment (Kumar *et al.* 2016; Palanikumar *et al.* 2025). Indigenous knowledge systems are embedded within local languages, social structures, value systems, institutions and customary laws and are largely based on experiential learning and naturalistic worldviews that differ significantly from formal scientific knowledge systems (International Union for Conservation of Nature 1997).

Human civilization has progressed from the Stone Age to the modern technological era through continuous observation, experimentation and adaptation. Agriculture and animal domestication form the foundation of early human societies, wherein communities gradually identified, domesticated and improved crops and livestock to meet subsistence requirements. Over successive generations, tribal communities refined ethno-agricultural and ethnoveterinary practices through trial and error, guided by intimate interactions with their local ecological conditions and resource availability (Kumar *et al.* 2012; Patel *et al.* 2018).

Ethno-agricultural and ethnoveterinary practices play a vital role in the conservation of plant and animal genetic resources, which are essential for ecological balance and long-term sustainability (Banerjee *et al.* 2014). These knowledge systems comprise locally evolved perceptions, information, and practices that enable tribal communities to manage land, crops, livestock and natural resources to fulfil their needs related to food, shelter, health, spiritual well-being and economic security. Indigenous knowledge is location-specific, dynamic, adaptive and continuously evolving in response to ecological, socio-economic and political changes.

Despite their significance, many indigenous agricultural and ethnoveterinary practices remain inadequately documented, scientifically validated and integrated into formal agricultural development and extension systems. Rapid urbanization, modernization of agriculture, environmental degradation, and socio-economic transitions have posed serious threats to the continuity and transmission of traditional knowledge (Kumar *et al.* 2016). Therefore, systematic documentation, analysis, and promotion of indigenous practices are essential to preserve this valuable heritage and to enhance sustainable and climate-resilient farming systems.

Agriculture constitutes the primary livelihood of the tribal population in the Kalrayan Hills of Tamil Nadu. The region’s varied topography, altitude and agro-climatic conditions support a diverse range of agricultural and horticultural crops, along with indigenous livestock

species. The Kalrayan Hills represent one of the prominent regions in the state where ethnographic and ethnoveterinary practices continue to be widely practiced for crop production, animal healthcare and livelihood generation. In particular, Villupuram district is known for its rich repository of indigenous knowledge related to agriculture and animal husbandry (Bashir *et al.* 2015; Callaby *et al.* 2016).

## 2. Materials and methods

The present study was conducted in the Kalrayan Hills region of Villupuram district, Tamil Nadu, which is predominantly inhabited by tribal communities practicing traditional agriculture and livestock rearing. An ex-post facto research design was adopted, as the variables under investigation had already occurred and were beyond the control of the researcher. The study area was selected purposively due to the prevalence of indigenous agricultural and ethnoveterinary practices among tribal farmers.

A multistage sampling technique was employed for the selection of respondents. In the first stage, villages with a high concentration of tribal households were identified. In the subsequent stage, tribal farmers actively engaged in farming and livestock rearing were selected randomly from the identified villages. A total of 300 tribal farmers were selected as respondents for the study, ensuring adequate representation of the tribal farming population in the selected villages. The sample size was considered statistically sufficient for behavioural research studies to generate reliable and generalizable findings.

Data were collected using a well-structured and pre-tested interview schedule developed based on relevant literature and expert consultation. The schedule covered personal, socio-economic, psychological and communication characteristics of the respondents, along with their extent of adoption of indigenous agricultural and ethnoveterinary practices.

Adoption behaviour was measured by assessing the extent to which respondents practiced selected indigenous agricultural and ethnoveterinary techniques. Scores were assigned based on the level of adoption, and respondents were categorized into low, medium and high adoption groups using appropriate statistical measures such as mean and standard deviation. The collected data were coded, tabulated and analyzed using suitable statistical tools such as frequency, percentage, mean and standard deviation to draw meaningful inferences.

## 3. Results

The results of the study revealed that a majority of the tribal farmers exhibited a medium level of adoption of indigenous agricultural and ethnoveterinary practices, followed by high and low adoption categories (Chandrasekar *et al.* 2017; Patel *et al.* 2018). The continued reliance on traditional practices indicates their practical relevance, cultural acceptance and economic feasibility in tribal farming systems. Practice-wise analysis showed a high level of adoption in indigenous agricultural practices, particularly those related to soil fertility management, seed treatment, crop protection and post-harvest operations (Balamurugan *et al.* 2017; Patel *et al.* 2018). Practices such as application of green leaf manure and farmyard manure, incorporation of crop residues, sun drying of harvested produce, use of neem-based pest control measures and indigenous storage methods recorded higher adoption percentages. Similarly,

adoption of ethnoveterinary practices was observed for the treatment of common livestock ailments such as fever, wounds, digestive disorders, parasitic infestations and post-calving care. Indigenous remedies using locally available medicinal plants, household materials and traditional preparations were widely practiced by the respondents (Avhad *et al.* 2015; Raina *et al.* 2016).

Overall adoption categorization indicated that a substantial proportion of respondents belonged to the medium to high adoption groups, reflecting the continued prevalence of indigenous knowledge systems among tribal farmers in the Kalrayan Hills region (Table 1, Figure 1 and Figure 2).

S. No.	Agriculture Practices	No. of Re- spondents	Per cent
<b>I.</b>	<b>Paddy</b>		
1	Soaking seeds for 24 hours in water and covering with paddy straw and bamboo leaves for early sprout	215	71.66
2	Seed rate @ 20-25 kg per acre	226	75.33
3	Burning of farm waste and trash on the nursery for better germination	189	63.00
4	Summer ploughing	195	65.00
5	Applying of green leaf manure and FYM	251	83.66
6	Incorporating crop residue and leaves of a tree as a manure	258	86.00
7	Sun drying of harvested paddy for one or two days in the field it self	268	89.33
8	Threshing by hitting the paddy bundles with wooden blocks	261	87.00
9	Parboiling of paddy for improving the edible quality of the rice	248	82.66
10	Irrigating from the channels when the well completely dries up	236	78.66
11	Grounding of rice in a heavy weight wooden grinder (Urral)	263	87.66
12	Using stingy bugs against caseworm	164	54.66
13	Bradcasting the crushed neem leaves in the paddy to reduce insect attack	266	88.66
14	Coating of cow dung solution in paddy grains for protection of pest and diseases	269	89.66
15	Covering rat holes with mud	248	82.66
	Mean		<b>79.04</b>
<b>II.</b>	<b>Tapioca</b>		
16	Selecting a setts with shorter internodes for planting	268	89.33
17	Cultivating banana as a inter crop between the rows	123	41.00
18	Application of pig manure for increased tuber size	222	74.00
19	Irrigating once in 15 days	257	85.66
20	Spraying of neem oil mixed with soap solution to control the pest and diseases	242	80.66

S. No.	Agriculture Practices	No. of Respondents	Per cent
21	Tapioca is cultivated in bench terrace	256	85.33
22	Selecting disease- free setts for propagation	261	87.00
23	Planting the setts within three hours after cutting	248	82.66
24	About 6-8 cuttings of 20 cm are obtained from mature stem, leaving the top tender shoot and woody bottom	248	82.66
25	The setts are planting the setts vertically at one inch depth in the soil	246	82.00
26	Cultivating Dolicho sp (India Been) as a smoother/cover crop in between the rows as an inter-crop.	224	74.66
27	Storage setts are cut and sun dried for a week and stored with 16% of moisture content	218	72.66
28	Mixing jatropha leaves with hot water (100 °C)is used to control aphids and white flies in tapioca	190	63.33
	Mean		<b>76.99</b>
<b>III.</b>	<b>Cumbu</b>		
29	Spreading of cumbu ear heads circularly to a height of 1 foot and cattle threshed	178	59.33
30	Drying of cumbu until a metallic sound is produced	188	62.66
31	Storing the cumbu in earthen pots covered with and tied cloth.	257	85.66
32	Spreading of Nochi leaves over the storage container to control pest	258	86.00
33	Mixing of seed purpose cumbu with dried neem leaves	257	85.66
34	Springing turmeric powder and ash solution (2Kg of turmeric powder + 8 Kg of ash + 200 litre of water per acre) to control sucking pests like aphids, hoppers etc.,	162	54.00
35	Cumbu ear heads are sun dried for two days and stored without seed separation by building a storage structure called 'Kudhir'.	256	85.33
36	Soaking the cumbu seeds in common salt solution before sowing to secure good germination under adverse conditions	262	87.33
37	Soaking the cumbu seeds in cow urine for half-an-hour and sun drying them before sowing to control head smut and to induce drought tolerance.	256	85.33
38	Sprinkling boiled water in the next day and immersed in ordinary water for some time before sowing in the filed give better in the filed better germination.	248	82.66
39	Country plough is run at the early stage of cumbu crop to ensure optimum plant population.	194	64.66

S. No.	Agriculture Practices	No. of Re- spondents	Per cent
40	Sowing cumbu during the tamil months Vaikasi - Aani (May-June) to avoid shoot fly and stem borer.	215	71.66
41	Sowing cowpea as an intercrop in cumbu to minimize stem borer attack due to its repellent smell.	161	53.66
42	Sowing lab-lab as an intercrop to reduce stem borer damage in cumbu.	193	64.33
43	Pouring neem cake extract drop by drop on the cumbu shoot to control shoot borer.	192	64.00
44	Dusting ash on the infected leaves of cumbu to prevent the pest incidence.	222	74.00
45	Dusting ash at milking stage to control ear head bugs.	192	64.00
46	Growing coriander as a mixed crop in cumbu to control the parasitic weed (Strigalutea).	146	48.66
47	A red / yellow/ dark cloth is tied to a long pole and fixed in the centre of the field to scare away the crows.	268	89.33
48	Mixing cumbu seeds with ash to prevent storage pests.	272	90.66
49	Local varieties are cultivate in dry lands to avoid more water coinciding with the harvesting stage.	276	92.00
50	Treating the cumbu seed treated with cow urine at 1:10 ratio to enhance germination.	267	89.00
51	Clewing dried cumbu grain gives, metallic sound and dryness.	248	82.66
52	Pounding cumbu into course powdery form and consumed	238	79.33
53	Dusting Chula ash in pearl millet fields to control green leaf hoppers sitting on inner side of leaves.	148	49.33
54	Storing cumbu seeds by mixing with ash.	257	85.66
	Mean		<b>59.90</b>
<b>IV.</b>	<b>General practices agriculture</b>		
55	Tying of polythene sheets to scare away the birds	266	88.66
56	Dusting of ash to control the pest	262	87.33
57	Sheep penning	267	89.00
58	Fumigating in closed container for ripening of fruits	273	91.00
59	Using neem seed kernel to control pest	258	86.00
60	Broadcasting enriched silt in the fields	150	50.00
61	Using of green chille and garlic extract to control aphid and jassid	218	72.66
62	Using of mounds, ridges and raised beds to reduce root rot problem.	252	84.00
63	Using mixture of gypsum and sugar for rodent birds	258	86.00
64	Broadcasting of cooked rice with milk to attract birds	252	84.00

S. No.	Agriculture Practices	No. of Respondents	Per cent
65	Spraying or tobacco extract to kill pest in crops	183	61.00
66	Soil and water conservation by use of stone terracing	207	69.00
67	Allowing pigs into the paddy field to control the nut sedge.	221	73.66
	Mean		<b>65.56</b>
	<b>Ethno practices under cow animal husbandry</b>		
<b>I.</b>	<b>Cow - Foot and mouth disease (FMD)</b>		
68	Giving local liquor or wine	188	62.66
69	Rubbing of jaggery in the mouth	170	56.66
70	Applying salt solution inside the mouth and between the hooves of the animal	121	40.33
	Mean		<b>53.21</b>
<b>II.</b>	<b>Selection of breed and feeding</b>		
71	Selecting of indigenous breed	232	77.33
72	Feeding dry roughages such as straw and hay to calving cows	167	55.66
73	Feeding all types of fodder to cows	265	88.33
74	Giving drinking water adequately to the cattle	261	87.00
	Mean		<b>77.08</b>
<b>III.</b>	<b>Care and management of dairy and pregnant cow</b>		
75	Isolating a pregnant cows from the rest house	266	88.66
76	Stopping milking 50 to 60 days before expected date of calving	252	84.00
77	Feeding roughages to pregnant cows	247	82.33
	Mean		<b>84.99</b>
<b>IV.</b>	<b>Ulcer on neck of the bullock</b>		
78	Applying boiled and cooled edible oil is applied over the neck to control rashes	190	63.33
79	Applying powdered coal paste on the ulcer part to minimize the pain	147	49.00
	Mean		<b>56.16</b>
<b>V.</b>	<b>Respiratory tract infection</b>		
80	Mixing a leaves of Thulasi (Ocimumcanum), arusha (Adhatodavasica), ginger, pepper, jaggery with water to make decoction and feed 2-3 times daily	125	41.66
81	Quashing the fruits of Kantakari (Solanum surattense) are soaked in goat urine overnight and filtered and squeezing into few drops the nostril	118	39.33
	Mean		<b>52.16</b>
<b>VI.</b>	<b>Dropping of placenta</b>		
82	Giving or three seeds of vellaikoundumani given with boiled bajra to the animal for immediate delivery	114	38.00

S. No.	Agriculture Practices	No. of Re- spondents	Per cent
83	Giving bambusa leaves for feeding to easy release of placenta	116	38.66
	Mean		<b>38.33</b>
<b>VII.</b>	<b>Mastitis in dairy animals</b>		
84	Applying Gheekumari (Aloe vera) – 1 or 3 petals Haldi (Turmeric) powder – 50gm Chunna (Lime stone) – 10 gm are made it paste and apply over the udder thrice a day	122	40.66
<b>VIII.</b>	<b>Treatment for the dislocated / fractured part of cow</b>		
85	Applying mixture of honey and pure ghee in the featured part	188	61.66
86	Applying of perandai pulp on the fractured part	121	40.33
87	Applying of mixture of salt, jaggery and turmeric powder in the featured part	120	40.00
88	Applying two vilvam fruits of partially burnt and ground water and make as paste to apply in the featured part	108	36.00
89	Applying fenugreek seed paste and bandaged in dislocated part.	152	50.66
	Mean		<b>57.16</b>
	<b>Sheep and goats</b>		
<b>I.</b>	<b>Blue tongue disease</b>		
90	Smearing a banana fruits with sesame oil for feed to animals for 2 to 3 times	122	40.66
91	Feeding leaf pulp of Aloe vera 100gm has to be administered daily.	115	38.33
	Mean		<b>39.49</b>
<b>II.</b>	<b>Eradication of the ecto – parasite</b>		
92	Applying of tobacco powder and edible oil mixture over the entire body of the animal	171	57.00
<b>III.</b>	<b>Flatulence</b>		
93	Feeding a mixture of onion and aerial root of banyan tree to the animal before	118	39.33
94	Applying salt in the tongue of the animal feeding tuber plant with onion mixture	114	38.00
95	Feeding of suspension of edible oil (100g), water and kerosene oil to the animals	122	40.66
	Mean		<b>39.33</b>
<b>IV.</b>	<b>Skin diseases</b>		
96	Applying of used engine oil over the skin	114	38.00
<b>V.</b>	<b>Cold</b>		
97	Dropping of bhoyrognijuice in the nose	94	31.33
<b>VI.</b>	<b>Diarrhea</b>		

S. No.	Agriculture Practices	No. of Respondents	Per cent
98	Oral administration of charcoal powder	168	56.00
99	Feeding leaf extract hupai	147	49.00
100	Feeding 3kg of steamed varagu grains	118	39.33
	Mean		<b>48.11</b>
<b>VII.</b>	<b>Unsuccessful conception</b>		
101	Feeding 200 – 300 ml of castor oil	148	49.33
102	Feeding of banana leaf extract	122	40.66
	Mean		<b>44.99</b>
<b>VIII.</b>	<b>Post – calving care</b>		
103	Feeding of 1- 2 kg jaggery dissolved in water to the animal immediately after calving	175	58.33
<b>I.</b>	<b>Poultry disease management</b>		
104	Spreading crushed leaves of sithapal (Annonasquamosa) inside poultry nest and lice collected over the leaves can be disposed hygienically	215	71.66
105	Applying garlic, tulasi, neem leaves, seethapal seeds, haldi each 10-20 gm are grounded together and boiled in 250ml of neem oil over the surface of the body of 10-15 birds	218	72.66
	Mean		<b>72.16</b>
<b>I.</b>	<b>Constipation</b>		
106	Giving castor oil, raw in seed oil can be given for 1-2 days according to species and body weight of animal.	218	72.66
107	Giving a decoction of 100 g of haldi (turmeric rhizome) in a litre of water may be given once for 1-3 days to age old animals.	116	38.66
	Mean		<b>55.66</b>
	<b>General diseases of animal husbandary</b>		
108	Pressing slightly heated local sword in the tooth for toothache control	108	36.00
109	Feeding little amount of cumin seeds for the gastroenteritis problem	151	50.33
110	Feeding well-grounded neem leaves, flowers and bark well and the cows for deworming	120	40.00
111	Applying Caetus (Carnegieagiganta) fluid is on the eyelids to control common eye disease	115	38.33
112	Giving salt mixed water control in digestion ( tympany)	106	35.33
113	Feeding tea waste powder in case of blood in urine	103	34.33
114	Applying turmeric paste against the fracture area	120	40.00
115	Pasting lime, garlic and turmeric paste to control open wounds	148	49.33
116	Pasting neem paste to control wounds of the animals	120	40.00
117	Applying ghee in case of crack of udder	118	39.33

S. No.	Agriculture Practices	No. of Re- spondents	Per cent
118	Applying Doorva ( <i>Calendula dactylon</i> Linn.) paste for bleeding of blood from any injury	103	34.33
119	Smearing of powder of Calamus ( <i>Acorus calamus</i> ) and the leaf extract of tulasi ( <i>Ocimum sanctum</i> ) mix on the body of animal prevent like and bovine flies	108	36.00
	Mean		<b>39.44</b>

Table 1: Distribution of respondents according to their practice wise adoption on recommended ethno agricultural and veterinary practices




Figure 1: Distribution of respondents according to their practice wise overall adoption level of respondents on ethno agricultural and veterinary practice

## 4. Discussion

The predominance of medium to high adoption levels among tribal farmers highlights the continued relevance of indigenous agricultural and ethnoveterinary practices in tribal livelihood systems. These findings suggest that traditional practices remain deeply embedded in the cultural and farming traditions of the study area. Higher adoption of indigenous agricultural practices may be attributed to their cost-effectiveness, eco-friendliness, easy availability of local resources and minimal dependence on external inputs. The results also indicate that indigenous practices are well adapted to the local agro-climatic conditions of the Kalrayan Hills.

The preference for ethnoveterinary practices can be explained by limited access to modern veterinary services in remote tribal areas, along with the trust developed through generations of experiential learning. Indigenous remedies are often perceived as safer, affordable and culturally acceptable alternatives to modern veterinary medicines. Factors such as age, farming experience, inheritance of traditional knowledge, social participation and access to



Figure 2: Indigenous agricultural and ethnoveterinary practices

indigenous resources were found to influence adoption behaviour. Older and more experienced farmers exhibited higher adoption levels, which underscores the role of experiential knowledge and intergenerational transmission in sustaining indigenous practices. These findings are in agreement with earlier studies emphasizing the importance of traditional knowledge systems in tribal agriculture.

The study reinforces the need for systematic documentation, scientific validation and integration of indigenous practices into formal agricultural extension programmes to ensure their preservation and effective utilization in sustainable and climate-resilient farming systems.

## 5. Conclusion

The study concluded that indigenous agricultural and ethnoveterinary practices remain an integral component of the livelihood systems of tribal farmers in the Kalrayan Hills of Tamil Nadu. A substantial proportion of respondents demonstrated medium to high levels of adoption, highlighting the continued relevance of traditional knowledge in sustainable agriculture and livestock management. Despite increasing exposure to modern agricultural technologies, tribal farmers continue to rely on indigenous practices due to their cost-effectiveness, eco-friendliness and cultural compatibility.

The findings underscore the need for systematic documentation, scientific validation, and integration of indigenous agricultural and ethnoveterinary practices into formal agricultural extension and development programmes. Strengthening participatory extension approaches

and promoting knowledge-sharing platforms can enhance the preservation and effective utilization of indigenous knowledge systems. Such efforts would contribute to sustainable agricultural development, biodiversity conservation and improved livelihood security among tribal communities in the Kalrayan Hills of Tamil Nadu.

## References

- Avhad SR, Kadian KS, Verma AK, Kale RB (2015). “Entrepreneurial behaviour of dairy farmers in Ahmednagar district of Maharashtra, India.” *Agricultural Science Digest*, **35**(1), 56–59.
- Balamurugan P, Senthilkumar A, Murugesan S (2017). “An analysis on socio-economic profile of backyard poultry farmers in Theni district of Tamil Nadu.” *International Journal of Science, Environment and Technology*, **6**(6), 3513–3519.
- Banerjee S, Pal S, Saha S (2014). “Indigenous knowledge systems in agriculture and natural resource management.” *Indian Journal of Traditional Knowledge*, **13**(3), 498–505.
- Bashir BP, Rajkamal PJ, Reeja GP (2015). “Adoption of modern animal husbandry practices by tribal livestock farmers of Attappady block in Kerala.” *Journal of Agricultural Science*, **7**(1–2), 17–21.
- Callaby RP, Toye A, Jennings A, Van Wyk O, Hanotte O, Mbole-Kariuki MN, Bronsvort BMdC, Kruuk LEB, Thumbi SM, Coetzer JAW, Conradie IC, Woolhouse MEJ, Kiara H (2016). “Seroprevalence of respiratory viral pathogens of indigenous calves in western Kenya.” *Research in Veterinary Science*, **108**, 120–124. doi:10.1016/j.rvsc.2016.08.007.
- Chandrasekar GK, Satyanarayan K, Jagadeeswary V, Shree JS (2017). “Relationship between socio-economic and psychological factors of dairy farmers with days open – A study in rural Karnataka.” *International Journal of Pure and Applied Biological Sciences*, **5**(1), 171–177.
- International Union for Conservation of Nature (1997). *Indigenous Peoples and Sustainability: Cases and Actions*. IUCN, Gland, Switzerland.
- Kumar BR, Prasad K, Sundarambal P (2012). “Role performance and level of tribal women farmers in Meghalaya.” *Indian Journal of Extension Education*, **12**(1), 60–63.
- Kumar M, Gupta J, Radhakrishnan A, Singh M (2016). “Socio-economic status and role of livestock to improve livelihood of tribes of Jharkhand.” *Research Journal of Agricultural Sciences*, **6**, 1421–1425. Special issue.
- Kumari J, Dubey R, Bose DK, Gupta V (2018). “A study on socio-economic condition of Tharu tribes in Bahraich district of Uttar Pradesh, India.” *Journal of Applied and Natural Science*, **10**(3), 939–944.
- Palanikumar K, Rajendran S, Murugan P (2025). “Indigenous technical knowledge in agriculture and livestock management among tribal communities.” *Indian Journal of Traditional Knowledge*, **24**(1), 45–52.
- Patel NK, Ashwar BK, Rajput MB, Prajapati MV (2018). “Personal and socio-economic characteristics of commercial dairy farmers and their association with economics.” *International Journal of Agricultural Sciences*, **10**(11), 6187–6191.
- Raina V, Bhusan B, Bakshi P, Khajuria S (2016). “Entrepreneurial behaviour of dairy farmers.” *Journal of Animal Research*, **6**(5), 947–953.

Vivekanandan P (1994). "Indigenous pest control methods." In *Conference on Indigenous Science and Technology*. Bharathidasan University, Tiruchirappalli, Tamil Nadu.

**! Publication & Reviewer Details****Publication Information**

- **Submitted:** *05 February 2026*
  - **Accepted:** *02 March 2026*
  - **Published (Online):** *03 March 2026*
- 

**Reviewer Information**

- **Reviewer 1:**  
**Dr. Manobharathi K**  
*Assistant Professor*  
*Mother Teresa College of Agriculture*  
*Pudukkottai, Tamil Nadu*
- **Reviewer 2:**  
**Dr. Mathuabirami V**  
*Assistant Professor*  
*Kaveri University*

**i Disclaimer/Publisher's Note**

The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). The publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#), which permits non-commercial use, sharing, and reproduction in any medium, provided the original work is properly cited and no modifications or adaptations are made.

**Affiliation:**

Kaviya P\*  
Agricultural Extension  
Chidambaram  
Tamil Nadu India  
E-mail: [kaviyadeepan66@gmail.com](mailto:kaviyadeepan66@gmail.com)

Natarajan M  
Agricultural Extension  
Chidambaram  
Tamil Nadu India  
E-mail: [mnextension@gmail.com](mailto:mnextension@gmail.com)  
URL: [https://annamalaiuniversity.ac.in/faculty\\_view.php?id=08163&dc=H10](https://annamalaiuniversity.ac.in/faculty_view.php?id=08163&dc=H10)